

References

1. Kajdashev IP, Zhdan VM, Rasin MS, Savchenko LG, Degtjar' NI, Borzyh OA, Mormol' IA, Gerasimenko ND, Shlykova OA, Jakimishina LI. *Ukrainskyi kardiologichnyi zhurnal – Ukrainian Journal of Cardiology* 2010;1(suppl):.
2. Kajdashev IP, Savchenko LG, Rasin MS. i soavt. XV Congress of Therapeutist of Ukraine,2004.171–175.
3. Leonova MV. *Farmateka* 2003;12:42–46 (in Russ).
4. Alvarez R, Reguero JR, Batalla A, Iglesias-Cubero G, Cortina A, Alvarez V, Coto E. Angiotensin converting enzyme and angiotensin II receptor 1 polymorphisms-association with early coronary disease *Cardiovasc Res* 1998;40:375–379.
5. Benetos A, Gautier S, Ricard S, Topouchian J, Cambien F. Influence of angiotensin-converting enzyme and angiotensin II type 1 receptor gene polymorphisms on aortic stiffness in normotensive and hypertensive patients. *Circulation* 1996;94:698–703.
6. Bonnardeaux A, Davies E, Jeunemaitre X, Fery I, Charru A, Clauser E, Tiret L, Cambien F, Corvol P and Soubrier F. Angiotensin II type 1 receptor gene polymorphisms in human essential hypertension. *Hypertension* 1994;24:63–69.
7. Danser AH, Schunkert H. Renin-angiotensin system gene polymorphisms: potential mechanisms for their association with cardiovascular diseases. *Eur J Pharmacol* 2000;410:303–316.
8. de Denus S, Zakrzewski-Jakubiak M, Dubé MP, Bélanger F, Lepage S, Leblanc M-H, Gossard D. Effects of AGTR1 A1166C gene polymorphism in patients with heart failure treated with candesartan. *Ann Pharmacother* 2008;42(7):925–32.
9. Gainer JV, Hunley TE, Kon V, Nadeau JH, Muldowney JAS, Brown NJ. Angiotensin II type I receptor polymorphism in African Americans lower frequency of the C-1166 variant. *Biochem Mol Biol Int* 1997;43:227–231.
10. Lapierre AV, Arce ME, Lopez JR, Ciuffo GM. Angiotensin II type 1 receptor A1166C GENE polymorphism and essential hypertension in San Luis. *Biocell v.30 n.3 Mendoza ago./dic. 2006.*
11. Liu Y, Zhuoma C, Shan G, Cui C, Hou S, Qin W, Cai D, Gesang L, Xiao Z, Pingcuo Z, Zheng H, Wu Z, Zhou W, Qiu C. A1166C Polymorphism of the Angiotensin II Type 1 Receptor Gene and Essential Hypertension in Han, Tibetan and Yi Populations. *Hypertens Res* 2002;25:515–521.
12. Stankovic A, Zivkovic M, Djuric T, Alavantic D. Angiotensin II type 1 receptor gene polymorphism and essential hypertension in Serbian population. *Clin Chim Acta* 2003;327:181–185.
13. Takami S, Katsuya T, Rakugi H, Noriyuki S, Nakata Y, Kamitani A, Miki T, Higaki J, Ogihara T. Angiotensin II Type 1 receptor gene polymorphism is associated with increase of left ventricular mass but not with hypertension. *AJH* 1998;2:316–321.

14. van Geel PP, Pinto YM, Buikema H, van Gilst WH. Is the A1166C gene polymorphism of the Angiotensin II type 1 receptor involved in cardiovascular disease? *Eur Heart J* 1998;19:G13–G17.
15. van Geel PP, Pinto YM, Voors AA, Buikema H, Oosterga M, Crijns HJ, van Gilst WH. Angiotensin II type 1 receptor A1166C gene polymorphism is associated with an increased response to angiotensin II in human arteries. *Hypertension* 2000;35(3):717–721.