

References

1. Abdullaev RY. *Ukrainskyi Radiologichnyi Zhurnal – Ukrainian Journal of Radiology* 2001;6(3):270-272 (in Ukr.).
2. Vilkenhof U, Kruk I. *Medliteratura*, 2008.240 p. (in Russ.).
3. Dzemeshkevich CL, Stivenson LU. *GEOTAR Med.*, 2000.288 p.(in Russ.).
4. Ivaniv JA. *Sertseva nedostatnist – Heart Failure* 2010;2:17–23 (in Ukr.).
5. Kovalenko BM, Chervonopiskaya OM, Illyash MG, Starshova OS. *Ukrainskyi Cardiologichnyi Zhurnal – Ukrainian Journal of Cardiology* 2004;5:77–81 (in Ukr.).
6. Matova EA, Pavlenko SI, Svishchenko EP, Mishchenko LA. *Ukrainskyi Cardiologichnyi Zhurnal – Ukrainian Journal of Cardiology* 2011;6:53–58 (in Russ.).
7. Matova EA, Pavlenko SI, Svishchenko EP. *Ukrainskyi Cardiologichnyi Zhurnal – Ukrainian Journal of Cardiology* 2003;1:60-65 (in Russ.).
8. Rekomendatsii Asotsiatsii kardiologiv Ukrainy. *Ukrainskyi Cardiologichnyi Zhurnal – Ukrainian Journal of Cardiology* 2013;1:15–32 (in Ukr.).
9. Rukovodstvo po kardiologii. Morion, 2008. 444–480 p.(in Russ.).
10. Abhayaratna W, Seward JB, Appleton CP. Left Atrial Size Physiologic Determinants and Clinical Applications. *J. Am. Coll. Cardiol* 2006;47:2357–2363.
11. Agricola E, Oppizzi M, Pisani M. Ischemic mitral regurgitation: mechanisms and echocardiographic classification. *Eur. J. Echocardiogr* 2008;9:207–221.
12. Amigoni M, Meris A, Thune JJ. Mitral regurgitation in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both: prognostic significance and relation to ventricular size and function. *Eur. Heart J* 2007;28:326–333.
13. Bursi F, Enriquez-Sarano M, Nkomo VT, Jacobsen SJ, Weston SA, Meverden RA. Heart failure and death after myocardial infarction in the community: the emerging role of mitral regurgitation. *Circulation* 2005;111:295–301.
14. Del Colle S, Milan A, De Castro S. Left atrial function and arterial hypertension. *J. Hum. Hypertens* 2008;3:28–32.
15. Grigioni F, Enriquez-Sarano M, Zehr KJ, Bailey KR, Tajik AJ. Ischemic mitral regurgitation: long-term outcome and prognostic implications with quantitative Doppler assessment. *Circulation* 2001;103:1759–1764.
16. Guidelines for the echocardiographic assessment of the right heart in adults: A report from the American society of Echocardiography endorsed by the European association of Echocardiography, a registered branch of the European society of cardiology and the Canadian Society of Echocardiography. *J. AM Soc. Echocardiogr* 2010;23:685–713.

17. Quinones MA, Otto CM, Stoddard M, Waggoner A, Zoghbi WA. Recommendations for quantification of Doppler echocardiography: a report from the Doppler Quantification Task Force of the Nomenclature and Standards Committee of the American Society of Echocardiography. *J. Am. Soc. Echocardiogr* 2002;15:167–184.
18. Zoghbi WA, Enriquez-Sarano M, Foster E, Grayburn PA. Recommendations for evaluation of the severity of native valvular regurgitation with two-dimensional and Doppler echocardiography. *Journal of the American Society of Echocardiography* 2003;16:567–612.