

References

1. Dziak HV, Zharinov OI. Fibryliatsiia peredserd: [monohrafiia] K.: Chetverta khvyliia, 2011.190p (in Ukr).
2. Lypnytskyi TM, Denysiuk VO, Kozlovskiy VO. *Bukovynskiy medychniy visnyk. – Medical Journal of Bukovina* 2003;7(2):131–133 (in Ukr).
3. Kozlovskiy VO. *Visnyk VDMU. – Journal of Vinnitsa State Medical University* 2003;1.1:18–20 (in Ukr).
4. Romanova EN, Yshchuk VA, Sychev OS. *Arytmologiya – Arrhythmology* 2012;3:24–32 (in Russ).
5. Kovalenko VM, Lutai MI. Sertsevo-sudynni zakhvoriuvannia. Rekomendatsii z diahnostryky, profilaktyky ta likuvannia K.: MORION, 2011. 408p (in Ukr).
6. Nadarasa K, Williams MJ. Single high oral dose amiodarone for cardioversion of recent onset atrial fibrillation. *Heart Lung Circ.* 2012;21(8):444–448.
7. Kowey PR, Breithardt G, Camm J, Crijns H, Dorian P, Le Heuzey JY, Pedrazzini L, Prystowsky EN, Salette G, Schwartz PJ, Torp-Pedersen C, Weintraub W. Physician stated atrial fibrillation management in light of treatment guidelines: data from an international, observational prospective survey. *Clin. Cardiol.* 2010;33 (3):172–178.
8. Camm AJ, Lip GY, De Caterina R. 2012 focused update of the ESC Guidelines for the management of atrial fibrillation: an update of the 2010 ESC Guidelines for the management of atrial fibrillation. Developed with the special contribution of the European Heart Rhythm Association. *Eur Heart J.* 2012;33(21):2719–2747.