

References

1. Amosova EN, Yao Syuy, Bezrodnyiy AB, Andreev EV, Prudkiy IV, Shishkina NV, Gerula OM, Syichenko YuA, Gorda II, Katsitadze IYu. *Ukrayinskyi kardiologichnyi zhurnal – Ukr J Cardiol* 2012;6:62–67 (in Russ).
2. Amosova KM, Parkhomenko OM, Rudenko YuV, Andryeyev YeV, Shumakov VO. *Ukrayinskyi kardiologichnyi zhurnal – Ukr J Cardiol* 2003;2:134–139 (in Ukr).
3. Byul A, Tsefal P. *DiaSoft*,2002.608 p (in Russ).
4. Kovalenko VM, Nesukay OH, Danylenko OO. *Ukrayinskyi kardiologichnyi zhurnal – Ukr J Cardiol* 2013;6:XXXXX (in Ukr).
5. Kovalenko VN, Nesukay OH., Danylenko OO. *Sertse i sudyny – Heart and vessels* 2013;4:XXXX (in Ukr).
6. Kovalenko VM, Nesukay OH, Danylenko OO, Polenova NS, Titov YeYu. *Ukrayinskyi medychnyi chasopys – Ukr Med J* 2013;2(94):183–187 (in Ukr).
7. Kovalenko VM, Nesukay OH, Polenova NS, Titov YeYu, Danylenko OO. *Ukrayinskyi kardiologichnyi zhurnal – Ukr J Cardiol* 2012;6:103–109 (in Ukr).
8. Plehanov AV. SPb.:Izd-vo SPbGUEF,2010.96 p (in Russ).
9. Shumakov VO, Talayeva TV, Tereshkevych LP, Malynovs'ka IE. *Ukrayinskyi kardiologichnyi zhurnal. – Ukr J Cardiol* 2011;4:25–30 (in Ukr).
10. Hanley JA, McNeil BJ. A method of comparing area under ROC - curves derived from the same cases. *Radiology* 1983;148(3):839–843.
11. Krone RJ, Gillespie JA, Weld FM, Miller JP, Moss AJ. Low-level exercise testing after myocardial infarction: usefulness in enhancing clinical risk stratification. *Circulation* 1985;71:80–89.
12. Kymälä MM, Antila M, Kivistö SM. Can strain rate imaging predict recovery of contraction after acute myocardial infarction? *Eur J Echocardiogr* 2011;12(5):364–371.
13. Leitman M, Lysyansky P, Sidenko S, Shir V, Peleg E, Binenbaum M, Kaluski E, Krakover R, Vered Z. Two-dimensional strain – a novel software for real-time quantitative echocardiographic assessment of myocardial function. *J Am Soc Echocardiogr* 2004;17(10):1021–1029.
14. Nucifora G, Bertini M, Ajmone Marsan N, Scholte AJ, Siebelink HM, Holman ER, Schalij MJ, van der Wall EE, Bax JJ, Delgado V. Temporal evolution of left ventricular dyssynchrony after myocardial infarction: relation with changes in left ventricular systolic function. *Eur Heart J Cardiovasc Imaging* 2012;13(12):1041–1046.
15. Nucifora G, Bertini M, Marsan NA, Scholte AJ, Siebelink HM, Holman ER, Schalij MJ, van der Wall EE, Bax JJ, Delgado V. Impact of left ventricular dyssynchrony early on left ventricular function after first acute myocardial infarction. *Am J Cardiol* 2010;105(3):306–311.

16. Sachdev V, Aletras AH, Padmanabhan S. Myocardial strain decreases with increasing transmural extent of infarction: a Doppler echocardiographic and magnetic resonance correlation study. *J Am Soc Echocardiogr* 2006;19(1):34–39.
17. Senaratne MP, Hsu LA, Rossall RE, Kappagoda CT. Exercise testing after myocardial infarction: relative values of the low level pre-discharge and the post-discharge exercise test. *J Am Coll Cardiol* 1988;12(6):1416–1422.
18. Stanton T, Leano R, Marwick TH. Prediction of all-cause mortality from global longitudinal speckle strain: comparison with ejection fraction and wall motion scoring. *Circ Cardiovasc Imaging* 2009;2(5):356–364.
19. Stevenson R, Umachandran V, Ranjadayalan K, Wilkinson P, Marchant B, Timmis AD. Reassessment of treadmill stress testing for risk stratification in patients with acute myocardial infarction treated by thrombolysis. *Br Heart J* 1993;70:415–420.
20. Vartdal T, Brunvand H, Pettersen E. Early prediction of infarct size by strain Doppler echocardiography after coronary reperfusion. *J Am Coll Cardiol* 2007;49(16):1715–1721.
21. Vilella M, Vilella A, Barbera S, Franzosi MG, Maggioni AP. Prognostic significance of double product and inadequate double product response to maximal symptom-limited exercise stress-testing after myocardial infarction in 6296 patients treated with thrombolytic agents. GISSI-2 Investigators. *Amer Heart J* 1999;137:443–522.