

References

1. Popov VV, Bulanova NA, Ivanov GG. Modern target of antihypertensive therapy. Data from clinical studies. Part 1. *Rat Pharmacother Cardiol* 2012;8(1):88–94.
2. Syrenko YN Dihydropyridine calcium antagonists – a new reserve to reduce the risk of complications of hypertension. *Arterial hypertens* 2010;1:47–52.
3. Syrenko M. Hypertensive disease and Arterial hypertension: monohrafyya. Donetsk: Publisher Zaslavsky A.2011.304 p.
4. Timofeeva AV, Goryunova LE, Khaspekov GL, Kovalevsky DA, Skamrov AV, Bulkin OS, Talitskii KA Karpov YA, Bibilashvili RSh. Pharmacogenetics, pharmacogenomics in light of the problems associated with essential arterial hypertension. *Kardiologichesky vestnik.* 2007;2(1):5–12.
5. Tseluyko VY, Yakovleva L., Strokova SA. Influence of adherence to therapy on prognosis in patients after myocardial infarction. *Liky of Ukraine* 2012;1:40–43.
6. ACCORD Study Group, Cushman WC, Evans GW, Byington RP, Goff DC Jr, Grimm RH Jr, Cutler JA, Simons-Morton DG, Basile JN, Corson MA, Probstfield JL, Katz L, Peterson KA, Friedewald WT, Buse JB, Bigger JT, Gerstein HC, Ismail-Beigi F. Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med* 2010;362:1575–85.
7. Brugts JJ, Simoons ML Genetic influences of angiotensin-converting enzyme inhibitor response: an opportunity for personalizing therapy? *Expert Rev Cardiovasc Ther* 2012;10(8):1001-9. doi: 10.1586/erc.12.83.
8. Butler MG. Genetics of hypertension. Current status. *J Med Liban* 2010;58(3):175–8.
9. Dahlöf B, Sever PS, Poulter NR, Wedel H, Beevers DG, Caulfield M, Collins R, Kjeldsen SE, Kristinsson A, McInnes GT, Mehlsen J, Nieminen M, O'Brien E, Ostergren J; ASCOT Investigators. Prevention of cardiovascular events with an antihypertensive regimen of amlodipine adding perindopril as required versus atenolol adding bendroflumethiazide as required, in the Anglo-Scandinavian Cardiac Outcomes Trial) Blood Pressure Lowering Arm (ASCOT-BPLA): a multicentre randomised controlled trial. *Lancet* 2005;366:895–906.
10. De la Sierra A, Roca-Cusachs A, Redón J, Marín R, Luque M, de la Figuera M, García-García M, Falkon L; CENIT Study Working Group. Effectiveness and tolerability of fixed-dose combination enalapril plus nitrendipine in hypertensive patients: results of the 3-month observational, post-marketing, multicentre, prospective CENIT study. *Clin Drug Investig* 2009;29:459–69.
11. Elliot WJ, Meyer PM. Incident diabetes in clinical trials of antihypertensive drugs: a network meta-analysis. *Lancet* 2007;369:201–7.
12. Forette F, Seux ML, Staessen JA, Thijs L, Birkenhäger WH, Babarskiene MR, Babeanu S, Bossini A, Gil-Extremera B, Girerd X, Laks T, Lilov E, Moisseyev V, Tuomilehto J, Vanhanen H, Webster J, Yodfat Y, Fagard R. Prevention of dementia in randomised double-blind placebo-controlled Systolic Hypertension in Europe (Syst-Eur) trial. *Lancet* 1998;352:1347–51.

13. Forette F, Seux ML, Staessen JA, Thijs L, Babarskiene MR, Babeau S, Bossini A, Fagard R, Gil-Extremera B, Laks T, Kobalava Z, Sarti C, Tuomilehto J, Vanhanen H, Webster J, Yodfat Y, Birkenhäger WH; Systolic Hypertension in Europe Investigators. The prevention of dementia with antihypertensive treatment: new evidence from the Systolic Hypertension in Europe (Syst-Eur) study. *Arch Intern Med* 2002;162:2046–52.
14. Grant RW, Devita NG, Singer DE, Meigs JB. Polypharmacy and medication adherence in patients with type 2 diabetes. *Diabetes Care* 2003;26(5):1408–12.
15. Hansson L, Zanchetti A, Carruthers SG, Dahlöf B, Elmfeldt D, Julius S, Ménard J, Rahn KH, Wedel H, Westerling S. Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. HOT Study Group. *Lancet* 1998;351(9118):1755–62.
16. Jin J, Sklar GE, Min Sen Oh V, Chuen Li S. Factors affecting therapeutic compliance: A review from the patient's perspective. *Ther Clin Risk Manag* 2008;4(1):269–86.
17. Lewington S, Clarke R, Qizilbash N., Peto R, Collins R. Prospective Studies Collaboration. Age specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002;360:1903–13.
18. Mancia G, Fagard R, Narkiewicz K, Redon J, Zanchetti A, Böhm M, Christiaens T, Cifkova R, De Backer G, Dominiczak A, Galderisi M, Grobbee DE, Jaarsma T, Kirchhof P, Kjeldsen SE, Laurent S, Manolis AJ, Nilsson PM, Ruilope LM, Schmieder RE, Sirnes PA, Sleight P, Viigimaa M, Waeber B, Zannad F, Redon J, Dominiczak A, Narkiewicz K, Nilsson PM, Burnier M, Viigimaa M, Ambrosioni E, Caufield M, Coca A, Olsen MH, Schmieder RE, Tsiofis C, van de Borne P, Zamorano JL, Achenbach S, Baumgartner H, Bax JJ, Bueno H, Dean V, Deaton C, Erol C, Fagard R, Ferrari R, Hasdai D, Hoes AW, Kirchhof P, Knuuti J, Kolh P, Lancellotti P, Linhart A, Nihoyannopoulos P, Piepoli MF, Ponikowski P, Sirnes PA, Tamargo JL, Tendera M, Torbicki A, Wijns W, Windecker S, Clement DL, Coca A, Gillebert TC, Tendera M, Rosei EA, Ambrosioni E, Anker SD, Bauersachs J, Hitij JB, Caulfield M, De Buyzere M, De Geest S, Derumeaux GA, Erdine S, Farsang C, Funck-Brentano C, Gerc V, Germano G, Gielen S, Haller H, Hoes AW, Jordan J, Kahan T, Komajda M, Lovic D, Mahrholdt H, Olsen MH, Ostergren J, Parati G, Perk J, Polonia J, Popescu BA, Reiner Z, Rydén L, Sirenko Y, Stanton A, Struijker-Boudier H, Tsiofis C, van de Borne P, Vlachopoulos C, Volpe M, Wood DA. 2013 ESH/ESC guidelines for the management of arterial hypertension: the Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Eur Heart J* 2013;34(28):2159–219. doi: 10.1093/eurheartj/eht151.
19. Marín-Iranzo R, de la Sierra-Iserte A, Roca-Cusachs A, Oliván-Martínez J, Redón-Mas J, Luque-Otero M, de la Figuera-von Wichman M, Pontes-García C, Delgadillo-Duarte J; grupo del estudio ENVIDA. Double blind study of the efficacy and safety of the fixed dose combination of enalapril 10 mg/nitrendipine 20 mg versus the increase of amlopidine dose in essential hypertensive patients not controlled with amlodipine 5 mg. *Rev Clin Esp* 2005;205(9):418–24.
20. Roca-Cusachs A, Torres F, Horas M, Ríos J, Calvo G, Delgadillo J, Terán M; Spanish Nitrendipine/Enalapril Collaborative Study Group. Nitrendipine and enalapril combination therapy in mild to moderate hypertension: assessment of dose-response relationship by a clinical trial of factorial design. *J Cardiovasc Pharmacol* 2001;38(6):840–9.