

References

1. Vatutin NT, Kalinkina NV, Shevelek AN. *Ukrainskyi kardiologichnyi zhurnal. – Ukr J Cardiol* 2010;1:46–50 (in Russ).
2. 2012 focused update of the ESC Guidelines for the management of atrial fibrillation. An update of the 2010 ESC Guidelines for the management of atrial fibrillation. *Eur Heart J* 2012;33:2719–47.
3. ACC/AHA/ASE 2005 Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Amer Soc Echocardiogr* 2005;18:1440–63.
4. Anderson ME, Al-Khatib SM, Roden DM, Califf RM. Cardiac repolarization: current knowledge, critical gaps, and new approaches to drug development and patient management. *Am Heart J* 2002;144:769–81.
5. Cao H, Wang X, Huang H, Ying SZ, Gu YW, Wang T, Huang CX. Omega-3 fatty acids in the prevention of atrial fibrillation recurrences after cardioversion: a meta-analysis of randomized controlled trials. *Intern Med* 2012;51:2503–8.
6. Coronel R, Wilms-Schopman FJ, Den Ruijter HM. Dietary n-3 fatty acids promote arrhythmias during acute regional myocardial ischemia in isolated pig hearts. *Cardiovasc Res* 2007;73:386–94.
7. Devereux RB, Alonso DR, Lutas EM, Gottlieb GJ, Campo E, Sachs I, Reichek N. Echocardiographic assessment of left ventricular hypertrophy: comparison to necropsy findings. *Am J Cardiol* 1986;57:450–8.
8. Grimsgaard S, Bønaa KH, Hansen JB, Myhre ES. Effects of highly purified eicosapentaenoic acid and docosahexaenoic acid on hemodynamics in humans. *Am J Clin Nutr* 1998;68:52–9.
9. Shen MJ, Choi EK, Tan AY, Lin SF, Fishbein MC, Chen LS, Chen PS. Neural mechanisms of atrial arrhythmias. *Nat Rev Cardiol* 2011;9:30–9.
10. Lorincz I, Szabó Z, Simkó J. Atrial fibrillation and the autonomous nervous system. *Orv Hetil* 2008;149:2019–28.
11. Macleod JC, Macknight AD, Rodrigo GC. The electrical and mechanical response of adult guinea pig and rat ventricular myocytes to omega-3 polyunsaturated fatty acids. *Eur J Pharmac* 1998;356:261–70.
12. Materazzoa C, Piottia P, Mantovania C, Miceli R, Villani F. Atrial fibrillation after non-cardiac surgery: P-wave characteristics and Holter monitoring in risk assessment. *Eur J Cardiothorac Surg* 2007;31:812–6.
13. Ninio DM, Murphy KJ, Howe PR, Saint DA. Dietary fish oil protects against stretch-induced vulnerability to atrial fibrillation in a rabbit model. *J Cardiovasc Electrophysiol* 2005;16:1–6.

14. O'Keefe JH, Abuissa H, Sastre A, Steinhaus DM, Harris WS. Effects of omega-3 fatty acids on resting heart rate, heart rate recovery after exercise, and heart rate variability in men with healed myocardial infarctions and depressed ejections fractions. *Am J Cardiol* 2006;97:27–30.
15. Rakowski H, Appleton C, Chan KL, Dumesnil JG, Honos G, Jue J, Koilpillai C, Lepage S, Martin RP, Mercier LA, O'Kelly B, Prieur T, Sanfilippo A, Sasson Z, Alvarez N, Pruitt R, Thompson C, Tomlinson C. Canadian consensus recommendation for the measurement and reporting of diastolic dysfunction in echocardiography. *J Am Soc Echocardiogr* 1996;9:736–60.
16. Szabó Z, Kakuk G, Fülöp T. Effects of haemodialysis on maximum P wave duration and P wave dispersion. *Nephrol Dial Transplant* 2002;17:1634–8.
17. Varounis C, Dargès N, Maounis T, Panagiotakos D, Cokkinos DV. Atrial premature complexes and heart rate have prognostic significance in 1-month atrial fibrillation recurrence after electrical cardioversion. *Europace* 2007;9:633–7.
18. Vincenti A, Brambilla R, Fumagalli MG, Merola R, Pedretti S. Onset mechanism of paroxysmal atrial fibrillation detected by ambulatory Holter monitoring. *Europace* 2006;8:204–10.