

## References

1. Sychov OS, Kovalenko VM, Dzyak GV, Korkushko AV, Bobrov VO, Gryn VK, Zharinov OI, Tashuk VK, Yabluchanskii MI, Stychinskii OS, Yagenskii OV, Karpenko UV, Pravosudovich SO, Solovian GM, Frolov OI, Kuts VO, Romanova OM, Getman TV, Volkov DE, Cherednichenko MZ, Sribnaya OV, Zalevskii VP, Kravchuk BB, Lysohub SV, Mogilnitskii EV, Levchuk NP, Levchuk OV, Borodai AO, Shabilianova LO, Lobko OO. Diagnostic and management of atrial fibrillation. Recommendation of working group of the association of cardiologists of Ukraine. Kyiv, 2011.159 p. (in Ukr.) – [http://strazhesko.org.ua/inc/materials/Book\\_ukr\\_%B10.2011%5D\\_preview.pdf](http://strazhesko.org.ua/inc/materials/Book_ukr_%B10.2011%5D_preview.pdf)
2. Khalafian AA. STATISTICA 6. ООО «Binom-Press», 2010.528p (in Russ.).
3. Blackshear JL, Odell JA. Appendage obliteration to reduce stroke in cardiac surgical patients with atrial fibrillation. *Ann Thorac Surg* 1996;61:755–9.
4. Florkowski ChM. Sensitivity, Specificity, Receiver-Operating Characteristic (ROC) Curves and Likelihood Ratios: Communicating the Performance of Diagnostic Tests *Clin Biochem Rev.* 2008 August;29(Suppl 1):S83–S87.
5. Go AS, Hylek EM, Phillips KA, Chang Y, Henault LE, Selby JV et al. Prevalence of diagnosed atrial fibrillation in adults: national implications for rhythm management and stroke prevention: the AnTicoagulation and Risk Factors in Atrial Fibrillation (ATRIA) Study. *JAMA* 2001;285:2370–5.
6. Thom T, Haase N, Rosamond W, Howard VJ, Rumsfeld J, Manolio T. Heart disease and stroke statistics-2006 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 2006;113:e85–151.
7. Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke* 1991;22:983–8.