

References

1. Belenkov YuN. *Cardiologiya – Cardiology* 1996;1:56-59 (in Russ).
2. Zheleznyak IS. Sankt-Peterburg, 2009.122p. (in Russ)
3. Lutai MI, Nemchina EA, Tsyzh AV, Golikova IP. *Ukrainskii kardiologichnyi zhurnal – Ukr Cardiol J* 2008;4:23–6 (in Russ).
4. Stukalova OV, Sinitsin VE, Ternovoy SK. *Meditsinskaya vizualizatsia – Medical visualization* 2004;5:18–23 (in Russ).
5. Trufanov HE, Zheleznyak IS, Rud SD, Menkov IA. ELBI, 2012.64p (in Russ).
6. Abdel-Aty H, Cocker M, Meek C, Tyberg JV, Friedrich MG. Edema as a very early marker for acute myocardial ischemia: a cardiovascular magnetic resonance study. *J Am Coll Cardiol* 2009; 53(14):1194–201
7. Abdel-Aty H, Zagrosek A, Schulz-Menger J, Taylor AJ, Messroghli D, Kumar A, Gross M, Dietz R, Friedrich MG. Delayed Enhancement and T2-Weighted Cardiovascular Magnetic Resonance Imaging Differentiate Acute From Chronic Myocardial Infarction. *Circulation* 2004 109(20):2411–16.
8. Aletras AH, Tilak GS, Natanzon A, Hsu LY, Gonzalez FM, Hoyt RF Jr, Arai AE. Retrospective determination of the area at risk for reperfused acute myocardial infarction with T2-weighted cardiac magnetic resonance imaging: histopathological and displacement encoding with stimulated echoes (DENSE) functional validations. *Circulation* 2006; 113(15):1865–70.
9. Alexánderon E, Jácome R, Romero E, Peña-Cabral M, Meléndez G, Kimura-Hayama E, Meave A. The importance of multi-imaging diagnosis in cardiology. *Arch Cardiol Mex* 2011;81(2):154–7.
10. Amano Y, Tachi M, Tani H, Mizuno K, Kobayashi Y, Kumita S. T2-weighted cardiac magnetic resonance imaging of edema in myocardial diseases. *Sci World J* 2012;37(3):1865–70.
11. Antony R, Daghem M, McCann GP, Daghem S, Moon J, Pennell DJ, Neubauer S, Dargie HJ, Berry C, Payne J, Petrie MC, Hawkins NM. Cardiovascular magnetic resonance activity in the United Kingdom: a survey on behalf of the British Society of Cardiovascular Magnetic Resonance. *J Cardiovasc Magn Reson* 2011; 13:57–63.
12. Baer FM, Theissen P, Crnac J, Schmidt M, Deutsch HJ, Sechtem U, Schicha H, Erdmann E. Head to head comparison of dobutamine-transoesophageal echocardiography and dobutamine-magnetic resonance imaging for the prediction of left ventricular functional recovery in patients with chronic coronary artery disease. *Eur Heart J* 2000;21(12):981–91.
13. Bettencourt N, Chiribiri A, Schuster A, Nagel E. Assessment of myocardial ischemia and viability using cardiac magnetic resonance. *Curr Heart Fail Rep* 2009;6(3):142–53.
14. Chiribiri A, Ishida M, Nagel E, Botnar RM. Coronary imaging with cardiovascular magnetic resonance: current state of the art. *Prog Cardiovasc Dis* 2011;54(3):240–52.

15. Choi KM, Kim RJ, Gubernikoff G, Vargas JD, Parker M, Judd RM. Transmural extent of acute myocardial infarction predicts longterm improvement in contractile function. *Circulation* 2001;104(10):1101–17.
16. Coelho-Filho OR, Seabra LF, Mongeon FP, Abdullah SM, Francis SA, Blankstein R, Di Carli MF, Jerosch-Herold M, Kwong RY. Stress myocardial perfusion imaging by CMR provides strong prognostic value to cardiac events regardless of patient's sex. *JACC Cardiovasc Imaging* 2011;4(8):850–61.
17. De Certaines JD. Is biomedical nuclear magnetic resonance limited by a revisitable paradigm in physics? *Cell Mol Biol (Noisy-le-grand)*. 2005;51(7):655–61.
18. Eitel I, Desch S, Sareban M, Fuernau G, Gutberlet M, Schuler G, Thiele H. Prognostic significance and magnetic resonance imaging findings in aborted myocardial infarction after primary angioplasty. *Am Heart J* 2009; 158(5):806–13.
19. Eitel I, Friedrich MG. T2-weighted cardiovascular magnetic resonance in acute cardiac disease. *J Cardiovasc Magn Reson*. 2011;13:131–44.
20. Figtree GA, Lønborg J, Grieve SM, Ward MR, Bhindi R. Cardiac magnetic resonance imaging for the interventional cardiologist. *JACC Cardiovasc Interv* 2011;4(2):137–48.
21. Florian A, Jurcut R, Gingham C, Bogaert J. Cardiac magnetic resonance imaging in ischemic heart disease: a clinical review. *J Med Life* 2011; 4(4):330–45.
22. Franccone M, Bucciarelli-Ducci C, Carbone I, Canali E, Scardala R, Calabrese FA, Sardella G, Mancone M, Catalano C, Fedele F, Passariello R, Bogaert J, Agati L. Impact of primary coronary angioplasty delay on myocardial salvage, infarct size, and microvascular damage in patients with ST-segment elevation myocardial infarction: insight from cardiovascular magnetic resonance. *J Am Coll Cardiol* 2009;54(23):2145–53.
23. Glaveckaitė S, Valeviciene N, Laucevicius A, Celutkienė J, Rudys A, Tamosiūnas A. Cardiovascular magnetic resonance imaging for detection of myocardial viability in chronic ischemic left ventricular dysfunction. *Medicina (Kaunas)* 2009;45(8):585–99.
24. Greenwood JP, Maredia N, Younger JF, Brown JM, Nixon J, Everett CC, Bijsterveld P, Ridgway JP, Radjenovic A, Dickinson CJ, Ball SG, Plein S. Cardiovascular magnetic resonance and single-photon emission computed tomography for diagnosis of coronary heart disease (CE-MARC): a prospective trial. *Lancet* 2012; 379(9814):453–60.
25. Gratz S, Kaiser W, Höffken H. Diagnostic imaging in patients with coronary artery disease: the nuclear medicine physicians' view. *Dtsch Med Wochenschr* 2011;136(41):2094–9.
26. Heydari B, Jerosch-Herold M, Kwong RY. Assessment of myocardial ischemia with cardiovascular magnetic resonance. *Prog Cardiovasc Dis* 2011;54(3):191–203.
27. Hombach V, Merkle N, Rasche V, Bernhardt P, Rottbauer W. The role of cardiovascular magnetic resonance imaging in the diagnosis and prognosis of patients with heart failure. *Herz* 2011; 36(2):84–93.
28. Iacucci I., Carbone I., Cannavale G., Conti B., Iampieri I., Rosati R., Sardella G., Frustaci A., Fedele F., Catalano C., Franccone M. Myocardial oedema as the sole marker of acute injury in Takotsubo cardiomyopathy: a cardiovascular magnetic resonance (CMR) study. *Radiol Med* 2013; 25(4):234–41.

29. Kim RJ, Wu E, Rafael A, Chen EL, Parker MA, Simonetti O, Klocke FJ, Bonow RO, Judd RM. The use of contrast-enhanced magnetic resonance imaging to identify reversible myocardial dysfunction. *N Engl J Med* 2000; 343(20):1445–53.
30. Kim RJ, Fieno DS, Parrish TB, Harris K, Chen EL, Simonetti O, Bundy J, Finn JP, Klocke FJ, Judd RM. Relationship of MRI delayed contrast enhancement to irreversible injury, infarct age, and contractile function. *Circulation* 1999;100(19):1992–2002.
31. Klein HM, Meyners W, Neeb B, Labenz J, Truümmler KH. Cardiac magnetic resonance imaging using an open 0.35 T system. *J Comput Assist Tomogr* 2007;31(3):430–4.
32. Klem I, Shah DJ, White RD, Pennell DJ, van Rossum AC, Regenfus M, Sechtem U, Schwartzman PR, Hunold P, Croisille P, Parker M, Judd RM, Kim RJ. Prognostic value of routine cardiac magnetic resonance assessment of left ventricular ejection fraction and myocardial damage: an international, multicenter study. *Circ Cardiovasc Imaging* 2011;4(6):610–619.
33. Kociemba A, Karmelita-Katulska K, Siniawski A, Lanocha M, Janus M, Stajgis M, Grajek S, Pyda M. Magnetic resonance imaging in the diagnostics of myocardial infarction. *Pol J Radiol* 2011;76(3):53–57.
34. Kumar A, Abdel-Aty H, Kriedemann I, Schulz-Menger J, Gross CM, Dietz R, Friedrich MG. Contrast-enhanced cardiovascular magnetic resonance imaging of right ventricular infarction. *J Am Coll Cardiol* 2006; 48(10):1969–76.
35. Kwong RY, Arai AE. Detecting patients with acute coronary syndrome in the chest pain center of the emergency department with cardiac magnetic resonance imaging. *Crit Pathw Cardiol* 2004;3(1):25–31.
36. Lockie T, Nagel E, Redwood S, Plein S. Use of cardiovascular magnetic resonance imaging in acute coronary syndromes. *Circulation* 2009;119(12):1671–81.
37. Macchia RJ, Termine JE, Buchen CD, Raymond V Damadian. Magnetic resonance imaging and the controversy of the 2003 Nobel Prize in Physiology or Medicine. *J Urol* 2007;178(3):783–785.
38. Mastouri R, Sawada SG, Mahenthiran J. Current noninvasive imaging techniques for detection of coronary artery disease. *Expert Rev Cardiovasc Ther* 2010;8(1):77–91.
39. Matthias G Friedrich, Han W Kim, Raymond J Kim. T2 weighted imaging to assess post-infarct myocardium at risk. *JACC Cardiovasc Imaging* 2011;4(9):1014–21.
40. Mavrogeni S, Bratis K, Kolovou G. Myocardial Ischemia and Viability by Cardiac Magnetic Resonance: The International Experience and the Greek Reality. *Hellenic J Cardiol* 2012;53(1):55–62.
41. Mollet NR, Dymarkowski S, Volders W, Wathiong J, Herbots L, Rademakers FE, Bogaert J. Visualization of ventricular thrombi with contrast-enhanced magnetic resonance imaging in patients with ischemic heart disease. *Circulation* 2002;106(23):2873–6.
42. Monmeneu JV, Bodí V, Sanchis J, López-Lereu MP, Mainar L, Núñez J, Chaustre F, Rumiz E, Chorro FJ, Llácer A. Cardiac magnetic resonance evaluation of edema after ST-elevation acute myocardial infarction. *Rev Esp Cardiol* 2009;62(8):858–66.

43. Monte GU, Drager LF, Souza FS, Avila LF, Parga Filho JR, César LA, Izaki M, Meneghetti JC, Rochitte CE, Kalil Filho R. Magnetic resonance vs technetium-99m pyrophosphate scintigraphy in the detection of perioperative myocardial necrosis. *Arq Bras Cardiol* 2008;91(2):113–8.
44. Oyama-Manabe N, Ishimori N, Sugimori H, Van Cauteren M, Kudo K, Manabe O, Okuaki T, Kamishima T, Ito YM, Tsutsui H, Tha KK, Terae S, Shirato H. Identification and further differentiation of subendocardial and transmural myocardial infarction by fast strain-encoded (SENC) magnetic resonance imaging at 3.0 Tesla. *Eur Radiol* 2011;21(11):2362–8.
45. Peel SA, Hussain T, Schaeffter T, Greil GF, Lagemaat MW, Botnar RM. Cross-sectional and in-plane coronary vessel wall imaging using a local inversion prepulse and spiral read-out: a comparison between 1.5 and 3 Tesla. *J Magn Reson Imaging* 2012;35(4):969–75.
46. Ripa RS, Nilsson JC, Wang Y, Søndergaard L, Jørgensen E, Kastrup J. Short- and long-term changes in myocardial function, morphology, edema, and infarct mass after ST-segment elevation myocardial infarction evaluated by serial magnetic resonance imaging. *Am Heart J* 2007;154(5):929–36.
47. Romero J, Xue X, Gonzalez W, Garcia MJ. CMR Imaging Assessing Viability in Patients With Chronic Ventricular Dysfunction Due to Coronary Artery Disease: A Meta-Analysis of Prospective Trials. *JACC Cardiovasc Imaging* 2012;5(5):494–508.
48. Ruder TD, Ebert LC, Khattab AA, Rieben R, Thali MJ, Kamat P. Edema is a sign of early acute myocardial infarction on post-mortem magnetic resonance imaging. *Forensic Sci Med Pathol* 2013;14(5):518–25.
49. Sem DS, Pellicchia M. NMR in the acceleration of drug discovery. *Curr Opin Drug Discov Devel* 2001;4(4):479–92.
50. So A, Lee TY. Quantitative myocardial CT perfusion: a pictorial review and the current state of technology development. *J Cardiovasc Comput Tomogr* 2011;5(6):467–81.
51. Vöhringer M, Flewitt JA, Green JD, Dharmakumar R, Wang J Jr, Tyberg JV, Friedrich MG. Oxygenation-sensitive CMR for assessing vasodilator-induced changes of myocardial oxygenation. *J Cardiovasc Magn Reson* 2010;12:20–31.
52. Wacker CM, Bock M, Hartlep AW, Beck G, van Kaick G, Ertl G, Bauer WR, Schad LR. Changes in myocardial oxygenation and perfusion under pharmacological stress with dipyridamole: assessment using T*2 and T1 measurements. *Magn Reson Med* 1999;41(4):686–95.
53. Wagner A, Mahrholdt H, Holly TA, Elliott MD, Regenfus M, Parker M, Klocke FJ, Bonow RO, Kim RJ, Judd RM. Contrast-enhanced MRI and routine single photon emission computed tomography (SPECT) perfusion imaging for detection of subendocardial myocardial infarcts: an imaging study. *Lancet* 2003;361:374–9.
54. Wright J, Bogaert J. Cardiac magnetic resonance imaging and ischaemic cardiomyopathies: what are the indications? *Presse Med* 2011;40:379–89
55. Yan AT, Shayne AJ, Brown KA, Gupta SN, Chan CW, Luu TM, Di Carli MF, Reynolds HG, Stevenson WG, Kwong RY. Characterization of the peri-infarct zone by contrast-enhanced cardiac magnetic resonance imaging is a powerful predictor of post-myocardial infarction mortality. *Circulation* 2006;114(1):32–9.