

References

1. Arablinskiy AV, Chechenov MH. Vozmozhnosti kompyuternoy tomografii i kompyuternoy tomograficheskoy angiografii v diagnostike vyrazhennosti zhirovoy infiltracii pecheni pri nealkogolnoy zhirovoy bolezni pecheni. *Radiologiya-praktika* 2010;2:10–20.
2. Lutay MI, Slobodskiy VA. Vzaimosvyaz mezhdru disfunkciey endoteliiya i tolerantnostyu k fizicheskoy nagruzke. *Ukr med chasopys* 2005;5:101–5.
3. Parhomenko AN, Lutay MYa. Novye podhody k gipolipidemicheskoy terapii u pacientov s ostrym infarktomyokarda. *Ukr med chasopys* 2005;4(90).
4. Bitar F, Lerman A, Akhter MW, Hatamizadeh P, Janmohamed M, Khan S, Elkayam U. Variable response of conductance and resistance coronary arteries to endothelial stimulation in patients with heart failure due to nonischemic dilated cardiomyopathy. *J Cardiovasc Pharmacol Ther* 2006;11:197–202.
5. Bugianesi E, Gastaldelli A. Hepatic and cardiac steatosis: Are they coupled? *Heart Fail Clin* 2012;8:663–70.
6. Charakida M, Masi S. Assessment of atherosclerosis: the role of flow-mediated dilatation. *Eur Heart J* 2005.31(23):2854–61.
7. Corretti MC, Celermajer D. Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J Am Coll Cardiol* 2002;39:257–65.
8. Fiorucci S, Mencarelli A, Palazzetti B, Del Soldato P, Morelli A, Ignarro LJ. An NO derivative of ursodeoxycholic acid protects against Fas-mediated liver injury by inhibiting caspase activity. *Proc Natl Acad Sci USA* 2001 Feb 27;98(5):2652–7.
9. Klosinska M, Rudzinski T, Grzelak P, Stefanczyk L, Drozd J, Krzeminska-Pakula M. Endothelium-dependent and -independent vasodilation is more attenuated in ischaemic than in non-ischaemic heart failure. *Eur J Heart Fail* 2009;11:765–70.
10. Kwan CY, Hsieh WT, To PN, Wang HD. New perspectives on vascular wall signaling: role of perivascular adipocytes and fibroblasts. *Acta Pharmacol Sin* 2010;31:1022–5.
11. Leeson P, Thorne S. Non-invasive measurement of endothelial function: effect on brachial artery dilatation of graded endothelial dependent and independent stimuli. *Heart* 1997;78:22–7.
12. Ma LN, Zhao SP, Gao M, Zhou QC, Fan P. Endothelial dysfunction associated with left ventricular diastolic dysfunction in patients with coronary heart disease. *Int Cardiol* 2000;72:275–9.
13. Papamichael CM, Aznaouridis KA. Effect of coffee on endothelial function in healthy subjects: the role of caffeine. *Clin Sci (Lond)* 2005;109:55–60.
14. Rivard AL, Steer CJ, Kren BT, Rodrigues CM, Castro RE, Bianco RW, Low WC. Administration of tauroursodeoxycholic acid (TUDCA) reduces apoptosis following myocardial infarction in rat. *Am J Chin Med* 2007;35(2):279–9.

15. Schindhelm RK, Diamant M. Liver alanine aminotransferase, insulin resistance and endothelial dysfunction in normotriglyceridaemic subjects with type 2 diabetes mellitus. *Eur J Clin Invest* 2005;35:369–74.
16. Shah A, Gkaliagkousi E, Ritter JM, Ferro A. Endothelial function and arterial compliance are not impaired in subjects with heart failure of non-ischemic origin. *J Card Fail* 2010;14:114–20.
17. Shechter M, Matetzky S, Arad M, Feinberg MS, Freimark D. Vascular endothelial function predicts mortality risk in patients with advanced ischaemic chronic heart failure. *Eur J Heart Fail* 2009;11:588–93.
18. Sorensen KE, Celermajer DS, Spiegelhalter DJ, Georgakopoulos D, Robinson J, Thomas O, Deanfield JE. Noninvasive measurement of human endothelium-dependent arterial responses: accuracy and reproducibility. *Brit Heart J* 1995;74:247–53.
19. Villanova N, Moscatiello S. Endothelial dysfunction and cardiovascular risk profile in nonalcoholic fatty liver disease. *Hepatology* 2005;42:473–80.