

References

1. Byul A, Tsefal P. DiaSoft, 2002. 608 p. (in Russ)
2. Vasiuk YuA, Hadzegova AB, Ivanova SV, Yushchuk EN, Amirbegishvili IM, Gerasimova NI. Tkanevaya dopplerografiya v ranney diagnostike funktsionalnykh narusheniy miokarda pri arterialnoy gipertenzii. *Ratsionalnaya farmakoterapiya v kardiologii – Rat Pharmacother in Cardiol* 2008;1:9–43. (in Russ)
3. Dzyak GV, Kolesnik MYu. Novyye vozmozhnosti v otsenke strukturno-funktsionalnogo sostoyaniya miokarda pri gipertonicheskoy bolezni. *Zdorovia Ukrainy – Health of Ukraine* 2013;1:24–25. (in Russ)
4. Zharinov OY, Kuts VO, Thor NV. Navantazhuvalni probi v kardiologii. Kiyiv: Medytsyna svitu 2006;89. (in Ukr)
5. Kovalenko VM, Nesukai OG, Danylenko OO, Titov EYu. Strukturno-funktsionalniy stan livogo shlunochka, vnutrishnoshlunochkova mehanichna dissinhroniya ta tolerantnist do flzichnogo navantazhennya u patsientiv, yaki perenesli infarkt miokarda iz zubtsem Q. *Ukrainskyi kardiologichnyi zhurnal – Ukr J Cardiol* 2014;1:13–22. (in Ukr)
6. Kovalenko VM, Nesukai OG, Polenova NS, Danylenko O, Titov EYu. Spekl-treking ehokardiografiya: normativni znachennya i rol metodu u vivchenni sistolichnoyi ta dlastolichnoyi funktsiyi livogo shlunochka. *Ukrainskyi kardiologichnyi zhurnal – Ukr J Cardiol* 2012;6:103–9. (in Ukr)
7. Nikitin NP, Kliland DF. Primenenie tkanevoy miokardialnoy dopplerehokardiographii v kardiologii. *Kardiologiya – Cardiology* 2002;3:66–79. (in Russ)
8. Bell SP, Nyland L, Tischler MD, McNabb M, Granzier H, LeWinter MM. Alterations in the determinants of diastolic suction during pacing tachycardia. *Circ Res* 2000;87(3):235–40.
9. Covell JW. Tissue structure and ventricular wall mechanics. *Circulation* 2008;118(7):699–701.
10. ESH/ESC guidelines for the management of arterial hypertension: the Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Eur Heart J* 2013;34(28):2159–219.
11. Goebel B, Gjesdal O, Kottke D. Regional and global myocardial function in patients with hypertensive heart disease: a two-dimensional ultrasound speckle tracking study. *Circulation* 2008;118:991–2.
12. Henein M, Mörner S, Lindmark K, Lindqvist P. Impaired left ventricular systolic function reserve limits cardiac output and exercise capacity in HFpEF patients due to systemic hypertension. *Int J Cardiol* 2013;168(2):1088–93.
13. Higginbotham MB, Morris KG, Williams RS, Coleman RE, Cobb FR. Physiologic basis for the age-related decline in aerobic work capacity. *Am J Cardiol.* 1986;57(15):1374–9.
14. Higginbotham MB, Morris KG, Williams RS, McHale PA, Coleman RE, Cobb FR. Regulation of stroke volume during submaximal and maximal upright exercise in normal man. *Circ Res.* 1986;58(2):281–91.

15. Ingels NB Jr. Myocardial fiber architecture and left ventricular function. *Technol Health Care*. 1997;5(1-2):45-52.
16. Kusunose K, Motoki H, Popovic ZB, Thomas JD, Klein AL, Marwick TH. Independent association of left atrial function with exercise capacity in patients with preserved ejection fraction. *Heart* 2012;98(17):1311-7.
17. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise J, Solomon S, Spencer KT, St John Sutton M, Stewart W. Recommendations for chamber quantification. *Eur J Echocardiogr* 2006;7(2):79-108.
18. Liu Y, Tsai WC, Lin C. Evidence of subtle left ventricular systolic dysfunction detected by automatic function imaging in patients with diastolic heart failure. Abstracts of EUROECHO the twelfth 2008;M 963:p.S120.
19. McCully RB, Roger VL, Ommen SR, Mahoney DW, Burger KN, Freeman WK, Pellikka PA. Outcomes of patients with reduced exercise capacity at time of exercise echocardiography. *Mayo Clin Proc* 2004;79(6):750-7.
20. Mor-Avi V, Lang RM, Badano LP, Belohlavek M, Cardim NM, Derumeaux G, Galderisi M, Marwick T, Nagueh SF, Sengupta PP, Sicari R, Smiseth OA, Smulevitz B, Takeuchi M, Thomas JD, Vannan M, Voigt JU, Zamorano JL. Current and evolving echocardiographic techniques for the quantitative evaluation of cardiac mechanics: ASE/EAE consensus statement on methodology and indications endorsed by the Japanese Society of Echocardiography. *Eur J Echocardiogr* 2011;12(3):167-205.
21. Mukherjee M, Dusaj R, Stewart K, Shapiro E, Katz R, Lewis J. Predictors of abnormal longitudinal strain using left ventricular speckle tracking in ambulatory hypertensive patients with preserved ejection fraction. Abstract. *J Am Coll Cardiol* 2013;61:E858.
22. Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, Waggoner AD, Flachskampf FA, Pellikka PA, Evangelisa A. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *Eur J Echocardiogr* 2009;10:165-93.
23. Niculescu N, Gherghinescu C, Cinteza M. Patients with Arterial Hypertension and no Coronary Heart Disease had Subclinical Left Ventricular Dysfunction as Assessed by Tissue Doppler Echocardiography. *Eur J of Echocardiogr* 2006;7:110.
24. Rademakers FE, Buchalter MB, Rogers WJ, Zerhouni EA, Weisfeldt ML, Weiss JL, Shapiro EP. Dissociation between left ventricular untwisting and filling. Accentuation by catecholamines. *Circulation* 1992;85(4):1572-81.