

References

1. Byul A, Tsefal P. DiaSoft,2002.608 p. (in Russ)
2. Dyadyik AI, Bagriy AE, Lebed IA, Yarovaya NF. Patogenez gipertrofii levogo zheludochka serdtsa u bolnyih arterialnyimi gipertoniyami. *Kardiologiya – Cardiology* 1995;1:59–63. (in Russ)
3. Kovalenko VM, Nesukay OG, Polenova NS Titov EYu, Danilenko OO. Spekl-treking ehokardiografiya: normativni znachennya i rol metodu u vivchenni sistolichnoyi ta dlastolichnoyi funktsiyi livogo shlunochka. *Ukrayinskyi kardiologichnyi zhurnal – Ukr J Cardiol* 2012;6:103–9. (in Ukr)
4. Saidova MA. Sovremennyye podhody k otsenke gipertrofii levogo zheludochka. Differentsialno-diagnosticheskie podhody. *Terapevticheskiy arkhiv – Ther Arch* 2012;84(4):5–11. (in Russ)
5. Aidietis A, Laucevicius A, Marinskis G. Hypertension and cardiac arrhythmias. *Curr Pharm Des* 2007;13:2545–55.
6. ESH/ESC guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Eur Heart J* 2013;34:2159–219.
7. Krumholz HV, Larson M, Levy D. Prognosis of left ventricular geometric patterns in the Framingham Heart Study. *J Am Coll Cardiol* 1995;25:879–84.
8. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise J, Solomon S, Spencer KT, St John Sutton M, Stewart W. Recommendations for chamber quantification. *Eur J Echocardiogr* 2006;7:79–108.
9. Mor-Avi V, Lang RM, Badano LP, Belohlavek M, Cardim NM, Derumeaux G, Galderisi M, Marwick T, Nagueh SF, Sengupta PP, Sicari R, Smiseth OA, Smulevitz B, Takeuchi M, Thomas JD, Vannan M, Voigt JU, Zamorano JL. Current and evolving echocardiographic techniques for the quantitative evaluation of cardiac mechanics: ASE/EAE consensus statement on methodology and indications endorsed by the Japanese society of echocardiography. *Eur J Echocardiogr* 2011;12:167–205.
10. Muiesan ML, Salvetti M, Monteduro C, Bonzi B, Paini A, Viola S, Poisa P, Rizzoni D, Castellano M, Agabiti-Rosei E. Left ventricular concentric geometry during treatment adversely affects cardiovascular prognosis in hypertensive patients. *Hypertens* 2004;43:731–8.
11. Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, Waggoner AD, Flachskampf FA, Pellikka PA, Evangelisa A. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *Eur J Echocardiogr* 2009;10:165–93.
12. Stefanadis C, Dernellis J, Toutouzas P. A clinical appraisal of left atrial function. *Eur Heart J* 2001;22:22–36.
13. Tan YT, Wenzelburger F, Lee E, Heatlie G, Leyva F, Patel K. et al. The pathophysiology of heart failure with normal ejection fraction: exercise echocardiography reveals complex abnormalities of both systolic and diastolic ventricular function involving torsion, untwist and longitudinal motion. *J Am Coll Cardiol* 2009;54:36–46.

14. Todaro MC, Choudhuri I, Belohlavek M, Jahangir A, Carerj S, Oreto L, Khandheria BK. New echocardiographic techniques for evaluation of left atrial mechanics. *Eur Heart J Cardiovasc Imaging* 2012;13:973–84.
15. Wang J, Khoury DS, Yue Y, Torre-Amione G, Nagueh SF. Preserved left ventricular twist and circumferential deformation, but depressed longitudinal and radial deformation in patients with diastolic heart failure. *Eur Heart J* 2008;29:1283–9.