

References

1. Bernick C, Kuller L, Dulberg C. et al. Cardiovascular Health Study Collaborative Research Group. Silent MRI infarcts and the risk of future stroke: the cardiovascular health study. *Neurology* 2001;57(7):1222–9.
2. Brant-Zawadzki M, Fein G, Van Dyke C, Kiernan R, Davenport L, de Groot J. MR imaging of the aging brain. *Am J Neuroradiol* 1985;6:675–82.
3. Brott T, Tomsick T, Feinberg W, Johnson C, Biller J, Broderick J, Kelly M, Frey J, Schwartz S, Blum C. Baseline silent cerebral infarction in the Asymptomatic Carotid Atherosclerosis Study. *Stroke* 1994;25:1122–9.
4. Camm AJ, Kirchhof P, Lip GY, Schotten U, Savelieva I, Ernst S, Van Gelder IC, Al-Attar N, Hindricks G, Prendergast B, Heidbuchel H, Alfieri O, Angelini A, Atar D, Colonna P, De Caterina R, De Sutter J, Goette A, Gorenek B, Heldal M, Hohloser SH, Kolh P, Le Heuzey JY, Ponikowski P, Rutten FH. Guidelines for the management of atrial fibrillation: The Task Force for the Management of Atrial Fibrillation of the European Society of Cardiology (ESC). *Eur Heart J* 2010;31(19):2369–429.
5. Ezekowitz MD, James KE, Nazarian SM, Davenport J, Broderick JP, Gupta SR, Thadani V, Meyer ML, Bridgers SL. for the Veterans Affairs Stroke Prevention in Nonrheumatic Atrial Fibrillation Investigators. Silent Cerebral Infarction in Patients With Nonrheumatic Atrial Fibrillation. *Circulation* 1995;92:2178–82.
6. Ferguson G, Eliasziw M, Barr H. Symptomatic Carotid Endarterectomy Trial Surgical Results in 1415 Patients for the North American Symptomatic Carotid Endarterectomy Trial (NASCET) Collaborators. *Stroke* 1999;30:1751–8.
7. Hara M, Ooie T, Yufu K, Tsunematsu Y, Kusakabe T, Ooga M, Saikawa T, Sakata T. Silent cortical strokes associated with atrial fibrillation. *Clin Cardiol* 1995;18:573–4.
8. Akoudad S, Sedaghat S, Hofman A, Koudstaal PJ, van der Lugt A, Ikram MA, Vernooij MW. Kidney function is related to cerebral small vessel disease. *Stroke* 2008;39(1):55–61.
9. Jeerakathil T, Wolf PA, Beiser A, Massaro J, Seshadri S, D'Agostino RB, DeCarli C. Stroke risk profile predicts white matter hyperintensity volume: the Framingham Study. *Stroke* 2004;35(8):1857–61.
10. Kleemann T, Becker T, Strauss M, Schneider S, Seidl K. Prevalence and clinical impact of left atrial thrombus and dense spontaneous echo contrast in patients with atrial fibrillation and low CHADS₂ score. *Eur J Echocardiogr* 2009;10(3):383–8.
11. Kobayashi A, Iguchi M, Shimizu S, Uchiyama S. Silent cerebral infarcts and cerebral white matter lesions in patients with nonvalvular atrial fibrillation. *J Stroke Cerebrovasc Dis* 2012;21(4):310–7.
12. Kutner M, Nachtsheim C, Neter J. Applied linear regression models. 4th ed. McGraw-Hill/Irwin, 2004.1396 p.

13. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise JS, Solomon SD, Spencer KT, Sutton MS, Stewart WJ. Recommendations for Chamber Quantification: A Report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, Developed in Conjunction with the European Association of Echocardiography, a Branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 2005;18:1440–63.
14. Mangla R, Kolar B, Almast J, Ekholm S. Border zone infarcts: pathophysiologic and imaging characteristics. *Radio Graphics* 2011;31:1201–14.
15. Pepi M, Evangelista A, Nihoyannopoulos P, Flachskampf FA, Athanassopoulos G, Colonna P, Habib G, Ringelstein EB, Sicari R, Zamorano JL, Sitges M, Caso P. Recommendations for echocardiography use in the diagnosis and management of cardiac sources of embolism European Association of Echocardiography (EAE) (a registered branch of the ESC). *Eur J Echocardiogr* 2010;11:461–76.
16. Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, Waggoner AD, Flachskampf FA, Pellikka PA, Evangelista A. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *J Am Soc Echocardiogr* 2009;22(2):107–33.
17. Pisters R, Olesen J, Lip G. The role of echocardiography in stroke risk assessment in patients with atrial fibrillation: is it additive or does it simply echo clinical risk factors? *Europace* 2012;14 (1):1–2.
18. Pollanen M, Deck J. The mechanism of embolic watershed infarction: experimental studies. *Can J Neurol Sci* 1990;17(4):395–8.
19. Price TR, Manolio TA, Kronmal RA, Kittner SJ, Yue NC, Robbins J, Anton-Culver H, O'Leary DH. Silent brain infarction on magnetic resonance imaging and neurological abnormalities in community-dwelling older adults: the Cardiovascular Health Study: CHS Collaborative Research Group. *Stroke* 1997;28:1158–64.
20. Russmann H, Vingerhoets F, Ghika J, Maeder P, Bogousslavsky J. Acute infarction limited to the lenticular nucleus: clinical, etiologic, and topographic features. *Arch Neurol* 2003;60 (3):351–5.
21. Sacco RL, Kasner SE, Broderick JP, Caplan LR, Connors JJ, Culebras A, Elkind MS, George MG, Hamdan AD, Higashida RT, Hoh BL, Janis LS, Kase CS, Kleindorfer DO, Lee JM, Moseley ME, Peterson ED, Turan TN, Valderrama AL, Vinters HV; An Updated Definition of Stroke for the 21st Century. A Statement for Healthcare Professionals From the American Heart Association/American Stroke Association. <http://stroke.ahajournals.org/content/early/2013/05/07/STR.0b013e318296aeca>.
22. Thambidorai SK, Murray RD, Parakh K, Shah TK, Black IW, Jasper SE, Li J, Apperson-Hansen C, Asher CR, Grimm RA, Klein AL. Utility of transesophageal echocardiography in identification of thrombogenic milieu in patients with atrial fibrillation (an ACUTE ancillary study). *Am J Cardiol* 2005;96:935–41.
23. Vermeer SE, Hollander M, van Dijk EJ, Hofman A, Koudstaal PJ, Breteler MM; Rotterdam Scan Study. Silent brain infarcts and white matter lesions increase stroke risk in the general population: the Rotterdam Scan Study. *Stroke* 2003;34:1126–9.

24. Vincelj J, Sokol I, Jaksic O. Prevalence and clinical significance of left atrial spontaneous echo contrast detected by transesophageal echocardiography. *Echocardiogr* 2002;19:319–24.
25. Vogels S., Emmelot-Vonk M., Verhaar H., Koek H. The association of chronic kidney disease with brain lesions on MRI or CT: a systematic review. *Maturitas* 2012;71 (4):331–6.
26. Wada M, Nagasawa H, Iseki C, Takahashi Y, Sato H, Arawaka S, Kawanami T, Kurita K, Daimon M, Kato T. Cerebral small vessel disease and chronic kidney disease (CKD): results of a cross-sectional study in community-based Japanese elderly. *J Neurol Sci* 2008;272(1–2):36–42.
27. Yonemura K, Kimura K, Minematsu K. Small centrum ovale infarcts on diffusion-weighted magnetic resonance imaging. *Stroke* 2002;33:1541–4.
28. Zabalgoitia M, Halperin JL, Pearce LA, Blackshear JL, Asinger RW, Hart RG. Transesophageal echocardiographic correlates of clinical risk of thromboembolism in nonvalvular atrial fibrillation. Stroke Prevention in Atrial Fibrillation III Investigators. *J Am Coll Cardiol* 1998;31 (7):1622–6.