

References

1. Ashcheulova TV, Kovalova OM, Saiied Mudzhakhid Abbas. *Ukrainskyi medychnyi almanakh – Ukrainian Medical Journal* 2013;16 (3):7–11.
2. Ashcheulova TV, Kovalova OM, Ambrosova TM, Saiied Mudzhakhid Abbas. *Ukrainskyi medychnyi almanakh – Ukrainian Medical Journal* 2012;15(5):11–4.
3. Kovaleva ON, Ashcheulova TV, Al Sheikkh Dyb KhKh. *Nauchnyie vedomosti Belhorodskogo gosudarstvennogo universiteta. Seriiia «Meditsina. Farmatsiia» – Scientific statements Belgorod State University. A series of "Medicine. Pharmacy"* 2012;22(141):20–3.
4. Colwell JA. Type 2 Diabetes, Pre-Diabetes, and the Metabolic Syndrome. *JAMA* 2011;306(2):215.
5. Bruun JM, Stallknecht B, Helge JW, Richelsen B. Interleukin-18 in plasma and adipose tissue: effects of obesity, insulin resistance, and weight loss. *Eur J Endocrinol* 2007;157:465–71.
6. van Exel E, Gussekloo J, de Craen AJ, Frölich M, Bootsma-Van Der Wiel A, Westendorp RG. Low production capacity of interleukin-10 associates with the metabolic syndrome and type 2 diabetes. The Leiden 85-plus study. *Diabetes* 2002;51:1088–92.
7. Mancia G, Laurent S, Agabiti-Rosei E, Ambrosioni E, Burnier M, Caulfield MJ, Cifkova R, Clément D, Coca A, Dominiczak A, Erdine S, Fagard R, Farsang C, Grassi G, Haller H, Heagerty A, Kjeldsen SE, Kiowski W, Mallion JM, Manolis A, Narkiewicz K, Nilsson P, Olsen MH, Rahn KH, Redon J, Rodicio J, Ruilope L, Schmieder RE, Struijker-Boudier HA, van Zwieten PA, Viigimaa M, Zanchetti A; Reappraisal of European guidelines on hypertension management: a European Society of Hypertension Task Force document. *J Hypertension* 2009;27:2121–58.
8. Troseid M, Seljeflot I, Hjerkin EM, Arnesen H. Interleukin-18 is a strong predictor of cardiovascular events in elderly men with metabolic syndrome. *Diabetes Care* 2009;32 (3):486–92.