

References

1. Diagnostica i likuvannia fibrillatsii predserdii. Recommendation of working group of the association of cardiologists of Ukraine. OS Sychov, VM Kovalenko, GV Dzyak, AV Korkushko, VO Bobrov, VK Gryn, OI Zharinov, VK Tashuk, MI Yabluchanskii, OS Stychinskii, OV Yagenskii, UV Karpenko, SO Pravosudovich, GM Solovian, O.I.Frolov, VO Kuts, OM Romanova, TV Getman, DE Volkov, MZ Cherednichenko, OV Sribnaya, VP Zalevskii, BB Kravchuk, SV Lysohub, EV Mogilnitskii, NP Levchuk, OV Levchuk, AO Borodai, LO Shablianova, OO Lobko. – Kyiv 2011.159 p.
2. Sychov OS, Borodai AA, Borodai ES. Silent cerebral infarcts: definition, clinical consequences prognosis and its association with atrial fibrillation. *Arrytmolohiia – Arrhythmology* 2014;3(11):26–37.
3. Sychov OS, Borodai AO, Fedkiv SV, Borodai ES, Getman TV, Rybak AYU. Predictors of silent cerebral infarctions of cardioembolic origin in patients with non-valvular atrial fibrillation. *Ukrainskyi kardiologichnyi zhurnal – Ukrainian cardiology journal* 2014;5:65–73.
4. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale. An updated literature review. *J Psychosom Res* 2002;52(2):69–77.
5. Dorian P, Jung W, Newman D, Paquette M, Wood K, Ayers GM, Camm J, Akhtar M, Luderitz B. The impairment of health-related quality of life in patients with intermittent atrial fibrillation: implications for the assessment of investigational therapy *J Am Coll Cardiol* 2000;36(4):1303–1309.
6. Ferguson G, Eliasziw M, Barr H, Clagett G, Barnes R, Wallace M, et al, Symptomatic Carotid Endarterectomy Trial Surgical Results in 1415 Patients for the North American Symptomatic Carotid Endarterectomy Trial (NASCET) Collaborators. *Stroke* 1999;30:1751–8.
7. Gehi AK, Sears S, Goli N, Walker TJ, Chung E, Schwartz J, Wood KA, Guise K, Mounsey JP. Psychopathology and symptoms of atrial fibrillation: implications for therapy. *Journal of Cardiovascular Electrophysiology*, 2012;23(5):473–478.
8. Shadi Kalantarian, Theodore A. Stern, MD; Moussa Mansour, MD; and Jeremy N. Ruskin, MD, Cognitive Impairment Associated With Atrial Fibrillation: A Meta-analysis *Ann Intern Med* 2013;158(5 Part 1):338–46.
9. Koteles F, Simor P. Modern health worries, somatosensory amplification and subjective symptoms: A longitudinal study. *Int J Behav Med* 2013;20:38–41.
10. Kupper N, Van den Broek KC, Widdershoven J, Denollet J. Subjectively reported symptoms in patients with persistent atrial fibrillation and emotional distress. *Front Psychol* 2013;4:1–9.
11. Kutner M, Nachtsheim C, Neter J. Applied Linear Regression Models, 4th ed. 2004

12. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise JS, Solomon SD, Spencer KT, Sutton MS, Stewart WJ; Recommendations for Chamber Quantification: A Report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, Developed in Conjunction with the European Association of Echocardiography, a Branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 2005;18:1440–63.
13. Marzona I, O'Donnell M, Teo K, Gao P, Anderson C, Bosch J, Yusuf S. Increased risk of cognitive and functional decline in patients with atrial fibrillation: results of the ONTARGET and TRANSCEND studies. *CMAJ* 2012;184(6):E329–36.
14. Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, Waggoner AD, Flachskampf FA, Pellikka PA, Evangelista A. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *J Am Soc Echocardiogr* 2009; 22(2):107–33.
15. Oldridge N, Höfer S, McGee H, Conroy R, Doyle F, Saner H; (for the HeartQoL Project Investigators). The HeartQoL: part II. Validation of a new core health-related quality of life questionnaire for patients with ischemic heart disease. *Eur J Prev Cardiol* 2014;21(1):98–106.
16. Russmann H, Vingerhoets F, Ghika J, Maeder P, Bogousslavsky J. Acute infarction limited to the lenticular nucleus: clinical, etiologic, and topographic features. *Arch Neurol* 2003 Mar; 60(3):351–5.
17. The EuroQol Group: Euro-Qol: A new facility for measurement of health-related quality of life. *Health Policy* 1990;16:199–208.
18. Thompson TS, Barksdale DJ, Sears SF, Mounsey JP, Pursell I, Gehi AK. The Effect of Anxiety and Depression on Symptoms Attributed to Atrial Fibrillation. *Pacing Clin Electrophysiol* 2014;37(4):439–46.
19. Thrall G, Lip GY, Carroll D, Lane D. Depression, anxiety, and quality of life in patients with atrial fibrillation. *Chest* 2007; 132:1259–64.
20. Udompanich S, Lip GY, Apostolakis S, Lane DA. Atrial fibrillation as a risk factor for cognitive impairment: a semi-systematic review. *QJM* 2013 Sep;106(9):795–802.
21. Vincelj J, Sokol I, Jaksic O. Prevalence and clinical significance of left atrial spontaneous echo contrast detected by transesophageal echocardiography. *Echocardiography* 2002;19:319–24.